



Knowledge grows

Cherry Crop Nutrition Soil Applied and Foliar Programme



Growth Stage	Bud Burst	Popcorn	Petal Fall	Stone Hardening	Developing Fruit	Post Harvest
YaraMila™	TROPICOTE 250 kg/ha applied as band		TROPICOTE 150 kg/ha applied as band			TROPICOTE 150-300 kg/ha applied as band
YaraVita™	ZINTRAC 700 1-2 l/ha		SENIPHOS 2-5 appl'ns of 10 l/ha at 7-14 day intervals			ZINTRAC 700 1-2 l/ha
YaraVita™	BUD BUILDER 5 - 10 l/ha		MAGFLO 300 2 x 0.5 l/ha at 10-14 days			BUD BUILDER 10 l/ha
YaraVita™		BORTRAC 150 1 l/ha		STOPIT 2 x appl'n of 10 l/ha 28 and 14 days before harvest		BORTRAC 150 1 l/ha
YaraVita™			MANTRAC PRO 2 x 1.0 l/ha at 10-14 days	FOLIAR POTASH 3 x Appl'n of 3-5 l/ha at 10-14 day intervals		FOLIAR POTASH 3-5 l/ha
YaraVita™	FERLEAF 100 1 l/ha		FERLEAF 100 1 l/ha repeated 3-8 times at 7-14 day intervals			COPTREL 0.5 l/ha
Analytical Tools	Water and Substrate Analysis - to identify limiting nutrients		Tissue Analysis - to identify limiting nutrients or deficiencies			

Read label before application of any YaraVita product

Soil applied products
 Foliar applied products
 Analytical tools

Yara Ireland Email: yaraireland@yara.com

Recommendations are based on Yara's field experience and field trial results

Disclaimer: The information contained herein is to the best of Yara's knowledge and belief, accurate.

Recommendations may need to be amended to ensure they comply with NVZ Nmax calculations where relevant.